

San Diego County AIDS Cases  
By Region\*  
Reported through March 31, 2000

	Cumulative Cases			Recent Cases**		
	Frequency	Percent	Rate/100,000	Frequency	Percent	Rate/100,000
Central San Diego	6677	64.8%	801.2	481	58.3%	57.7
East County	759	7.4%	149.1	59	7.2%	11.6
North County	1180	11.4%	128.8	107	13.0%	11.4
South Bay	798	7.7%	220.0	106	12.8%	29.2
Southeast San Diego	895	8.7%	426.2	72	8.7%	34.3
<b>TOTAL</b>	<b>10,309</b>	<b>100.0%</b>	<b>361.3</b>	<b>825</b>	<b>100.0%</b>	<b>28.9</b>

\*As defined by the Office of AIDS Coordination

\*\*Cases diagnosed during the period April 1, 1998 – March 31, 2000


Percent may not total 100 due to rounding

Division of Community Epidemiology, April, 2000


[MS-0399.001.016.003]

## San Diego County AIDS Cases Regional Proportion of Cases by Five Year Cohorts Reported through March 31, 2000


Central Region




East Region




North Region



South Bay Region



Southeast San Diego Region



Region: Central San Diego

	Countywide				Region: Central San Diego			
	Cumulative Cases		Recent Cases*		Cumulative Cases		Recent Cases*	
	No.	%	No.	%	No.	%	No.	%
<b>Mode of Transmission – Adults</b>								
Men having Sex with Men (MSM)	7834	76.4%	579	70.2%	5384	80.9%	356	74.0%
Injection Drug Use (IDU)	860	8.4%	112	13.6%	388	5.8%	54	11.2%
MSM+IDU	913	8.9%	67	8.1%	628	9.4%	49	10.2%
Hemophilia	57	0.6%	1	0.1%	21	0.3%	1	0.2%
Heterosexual	409	4.0%	57	6.9%	166	2.5%	18	3.7%
Transfusion	151	1.5%	4	0.5%	60	0.9%	1	0.2%
No Risk Specified/Other	33	0.3%	5	0.6%	10	0.2%	2	0.4%
<b>Mode of Transmission – Pediatric</b>								
Parent at risk/with HIV/AIDS	42	80.8%	0	0.0%	18	90.0%	0	0.0%
Pediatric Hemophilia	3	5.8%	0	0.0%	1	5.0%	0	0.0%
Pediatric Transfusion	7	13.5%	0	0.0%	1	5.0%	0	0.0%
<b>Race/Ethnicity</b>								
White	6891	66.8%	422	51.2%	4913	73.6%	287	59.7%
Black	1201	11.7%	120	14.5%	691	10.3%	76	15.8%
Hispanic	1970	19.1%	268	32.5%	945	14.2%	112	23.3%
Asian/Pacific Islander	191	1.9%	10	1.2%	91	1.4%	2	0.4%
Native American	56	0.5%	5	0.6%	37	0.6%	4	0.8%
<b>Gender</b>								
Male	9617	93.3%	750	90.9%	6389	95.7%	452	94.0%
Female	692	6.7%	75	9.1%	288	4.3%	29	6.0%
<b>Age Group</b>								
1-12	52	0.5%	0	0.0%	20	0.3%	0	0.0%
13-19	35	0.3%	3	0.4%	14	0.2%	1	0.2%
20-29	1810	17.6%	117	14.2%	1183	17.7%	65	13.5%
30-39	4828	46.8%	385	46.7%	3183	47.7%	229	47.6%
40-49	2571	24.9%	223	27.0%	1687	25.3%	131	27.2%
50+	1013	9.8%	97	11.7%	590	8.8%	55	11.5%
<b>TOTAL</b>	<b>10,309</b>	<b>100.0%</b>	<b>825</b>	<b>100.0%</b>	<b>6677</b>	<b>100.0%</b>	<b>481</b>	<b>100.0%</b>

\*Cases diagnosed during the period April 1, 1998 – March 31, 2000

Percent may not total 100 due to rounding

Division of Community Epidemiology, April, 2000

Region: Southeast San Diego

	Countywide				Region: Southeast San Diego			
	Cumulative Cases		Recent Cases		Cumulative Cases		Recent Cases	
	No.	%	No.	%	No.	%	No.	%
<b>Mode of Transmission – Adults</b>								
Men having Sex with Men (MSM)	7834	76.4%	579	70.2%	602	67.6%	42	58.3%
Injection Drug Use (IDU)	860	8.4%	112	13.6%	142	16.0%	19	26.4%
MSM+IDU	913	8.9%	67	8.1%	87	9.8%	5	6.9%
Hemophilia	57	0.6%	1	0.1%	3	0.3%	0	0.0%
Heterosexual	409	4.0%	57	6.9%	40	4.5%	6	8.3%
Transfusion	151	1.5%	4	0.5%	12	1.3%	0	0.0%
No Risk Specified/Other	33	0.3%	5	0.6%	4	0.4%	0	0.0%
<b>Mode of Transmission – Pediatric</b>								
Parent at risk/with HIV/AIDS	42	80.8%	0	0.0%	4	80.0%	0	0.0%
Pediatric Hemophilia	3	5.8%	0	0.0%	0	0.0%	0	0.0%
Pediatric Transfusion	7	13.5%	0	0.0%	1	20.0%	0	0.0%
<b>Race/Ethnicity</b>								
White	6891	66.8%	422	51.2%	355	39.7%	16	22.2%
Black	1201	11.7%	120	14.5%	263	29.4%	19	26.4%
Hispanic	1970	19.1%	268	32.5%	249	27.8%	34	47.2%
Asian/Pacific Islander	191	1.9%	10	1.2%	24	2.7%	3	4.2%
Native American	56	0.5%	5	0.6%	4	0.4%	0	0.0%
<b>Gender</b>								
Male	9617	93.3%	750	90.9%	805	89.9%	62	86.1%
Female	692	6.7%	75	9.1%	90	10.1%	10	13.9%
<b>Age Group</b>								
1-12	52	0.5%	0	0.0%	5	0.6%	0	0.0%
13-19	35	0.3%	3	0.4%	9	1.0%	0	0.0%
20-29	1810	17.6%	117	14.2%	172	19.2%	12	16.7%
30-39	4828	46.8%	385	46.7%	409	45.7%	26	36.1%
40-49	2571	24.9%	223	27.0%	215	24.0%	24	33.3%
50+	1013	9.8%	97	11.7%	85	9.5%	10	13.9%
<b>TOTAL</b>	<b>10,309</b>	<b>100.0%</b>	<b>825</b>	<b>100.0%</b>	<b>895</b>	<b>100.0%</b>	<b>72</b>	<b>100.0%</b>

\*Cases diagnosed during the period April 1, 1998 – March 31, 2000

Percent may not total 100 due to rounding

Division of Community Epidemiology, April, 2000

Region: North County

	Countywide				Region: North County			
	Cumulative Cases		Recent Cases*		Cumulative Cases		Recent Cases*	
	No.	%	No.	%	No.	%	No.	%
<b>Mode of Transmission – Adults</b>								
Men having Sex with Men (MSM)	7834	76.4%	579	70.2%	801	68.4%	69	64.5%
Injection Drug Use (IDU)	860	8.4%	112	13.6%	144	12.3%	17	15.9%
MSM+IDU	913	8.9%	67	8.1%	81	6.9%	7	6.5%
Hemophilia	57	0.6%	1	0.1%	12	1.0%	0	0.0%
Heterosexual	409	4.0%	57	6.9%	85	7.3%	13	12.1%
Transfusion	151	1.5%	4	0.5%	36	3.1%	1	0.9%
No Risk Specified/Other	33	0.3%	5	0.6%	12	1.0%	0	0.0%
<b>Mode of Transmission – Pediatric</b>								
Parent at risk/with HIV/AIDS	42	80.8%	0	0.0%	6	66.7%	0	0.0%
Pediatric Hemophilia	3	5.8%	0	0.0%	0	0.0%	0	0.0%
Pediatric Transfusion	7	13.5%	0	0.0%	3	33.3%	0	0.0%
<b>Race/Ethnicity</b>								
White	6891	66.8%	422	51.2%	820	69.5%	68	63.6%
Black	1201	11.7%	120	14.5%	81	6.9%	4	3.7%
Hispanic	1970	19.1%	268	32.5%	238	20.2%	31	29.0%
Asian/Pacific Islander	191	1.9%	10	1.2%	34	2.9%	4	3.7%
Native American	56	0.5%	5	0.6%	7	0.6%	0	0.0%
<b>Gender</b>								
Male	9617	93.3%	750	90.9%	1032	87.5%	90	84.1%
Female	692	6.7%	75	9.1%	148	12.5%	17	15.9%
<b>Age Group</b>								
1-12	52	0.5%	0	0.0%	9	0.8%	0	0.0%
13-19	35	0.3%	3	0.4%	5	0.4%	1	0.9%
20-29	1810	17.6%	117	14.2%	181	15.3%	10	9.3%
30-39	4828	46.8%	385	46.7%	511	43.3%	47	43.9%
40-49	2571	24.9%	223	27.0%	316	26.8%	33	30.8%
50+	1013	9.8%	97	11.7%	158	13.4%	16	14.9%
<b>TOTAL</b>	<b>10,309</b>	<b>100.0%</b>	<b>825</b>	<b>100.0%</b>	<b>1180</b>	<b>100.0%</b>	<b>107</b>	<b>100.0%</b>

\*Cases diagnosed during the period April 1, 1998 – March 31, 2000

Percent may not total 100 due to rounding

Division of Community Epidemiology, April, 2000

Region: East County

	Countywide				Region: East County			
	Cumulative Cases		Recent Cases*		Cumulative Cases		Recent Cases*	
	No.	%	No.	%	No.	%	No.	%
<b>Mode of Transmission – Adults</b>								
Men having Sex with Men (MSM)	7834	76.4%	579	70.2%	517	68.5%	39	66.1%
Injection Drug Use (IDU)	860	8.4%	112	13.6%	85	11.3%	8	13.6%
MSM+IDU	913	8.9%	67	8.1%	55	7.3%	2	3.4%
Hemophilia	57	0.6%	1	0.1%	18	2.4%	0	0.0%
Heterosexual	409	4.0%	57	6.9%	55	7.3%	8	13.6%
Transfusion	151	1.5%	4	0.5%	21	2.8%	0	0.0%
No Risk Specified/Other	33	0.3%	5	0.6%	4	0.5%	2	3.4%
<b>Mode of Transmission – Pediatric</b>								
Parent at risk/with HIV/AIDS	42	80.8%	0	0.0%	3	75.0%	0	0.0%
Pediatric Hemophilia	3	5.8%	0	0.0%	1	25.0%	0	0.0%
Pediatric Transfusion	7	13.5%	0	0.0%	0	0.0%	0	0.0%
<b>Race/Ethnicity</b>								
White	6891	66.8%	422	51.2%	546	71.9%	29	49.2%
Black	1201	11.7%	120	14.5%	76	10.0%	12	20.3%
Hispanic	1970	19.1%	268	32.5%	118	15.5%	16	27.1%
Asian/Pacific Islander	191	1.9%	10	1.2%	12	1.6%	1	1.7%
Native American	56	0.5%	5	0.6%	7	0.9%	1	1.7%
<b>Gender</b>								
Male	9617	93.3%	750	90.9%	684	90.1%	50	84.7%
Female	692	6.7%	75	9.1%	75	9.9%	9	15.3%
<b>Age Group</b>								
1-12	52	0.5%	0	0.0%	4	0.5%	0	0.0%
13-19	35	0.3%	3	0.4%	4	0.5%	1	1.7%
20-29	1810	17.6%	117	14.2%	117	15.4%	7	11.9%
30-39	4828	46.8%	385	46.7%	347	45.7%	29	49.2%
40-49	2571	24.9%	223	27.0%	192	25.3%	15	25.4%
50+	1013	9.8%	97	11.7%	95	12.5%	7	11.9%
<b>TOTAL</b>	<b>10,309</b>	<b>100.0%</b>	<b>825</b>	<b>100.0%</b>	<b>759</b>	<b>100.0%</b>	<b>59</b>	<b>100.0</b>

\*Cases diagnosed during the period April 1, 1998 – March 31, 2000

Percent may not total 100 due to rounding

Division of Community Epidemiology, April, 2000

Region: South Bay

	Countywide				Region: South Bay			
	Cumulative Cases		Recent Cases*		Cumulative Cases		Recent Cases*	
	No.	%	No.	%	No.	%	No.	%
<b>Mode of Transmission – Adults</b>								
Men having Sex with Men (MSM)	7834	76.4%	579	70.2%	530	67.6%	73	68.9%
Injection Drug Use (IDU)	860	8.4%	112	13.6%	101	12.9%	14	13.2%
MSM+IDU	913	8.9%	67	8.1%	62	7.9%	4	3.8%
Hemophilia	57	0.6%	1	0.1%	3	0.4%	0	0.0%
Heterosexual	409	4.0%	57	6.9%	63	8.0%	12	11.3%
Transfusion	151	1.5%	4	0.5%	22	2.8%	2	1.9%
No Risk Specified/Other	33	0.3%	5	0.6%	3	0.4%	1	0.9%
<b>Mode of Transmission – Pediatric</b>								
Parent at risk/with HIV/AIDS	42	80.8%	0	0.0%	11	78.6%	0	0.0%
Pediatric Hemophilia	3	5.8%	0	0.0%	1	7.1%	0	0.0%
Pediatric Transfusion	7	13.5%	0	0.0%	2	14.3%	0	0.0%
<b>Race/Ethnicity</b>								
White	6891	66.8%	422	51.2%	257	32.2%	22	20.8%
Black	1201	11.7%	120	14.5%	90	11.3%	9	8.5%
Hispanic	1970	19.1%	268	32.5%	420	52.6%	75	70.8%
Asian/Pacific Islander	191	1.9%	10	1.2%	30	3.8%	0	0.0%
Native American	56	0.5%	5	0.6%	1	0.1%	0	0.0%
<b>Gender</b>								
Male	9617	93.3%	750	90.9%	707	88.6%	96	90.6%
Female	692	6.7%	75	9.1%	91	11.4%	10	9.4%
<b>Age Group</b>								
1-12	52	0.5%	0	0.0%	14	1.8%	0	0.0%
13-19	35	0.3%	3	0.4%	3	0.4%	0	0.0%
20-29	1810	17.6%	117	14.2%	157	19.7%	23	21.7%
30-39	4828	46.8%	385	46.7%	378	47.4%	54	50.9%
40-49	2571	24.9%	223	27.0%	161	20.2%	20	18.9%
50+	1013	9.8%	97	11.7%	85	10.7%	9	8.4%
<b>TOTAL</b>	<b>10,309</b>	<b>100.0%</b>	<b>825</b>	<b>100.0%</b>	<b>798</b>	<b>100.0%</b>	<b>106</b>	<b>100.0%</b>

\*Cases diagnosed during the period April 1, 1998 – March 31, 2000

Percent may not total 100 due to rounding

Division of Community Epidemiology, April, 2000